



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 680 764 A2**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **95106821.2**

(51) Int. Cl.⁶: **A61K 38/48, C12N 9/96,
A61K 47/24, //(A61K38/48,
38:35)**

(22) Date of filing: **05.05.95**

(30) Priority: **06.05.94 DE 4416166**

(43) Date of publication of application:
08.11.95 Bulletin 95/45

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB IT LI NL SE

(71) Applicant: **IMMUNO Aktiengesellschaft
Industriestrasse 67
A-1221 Wien (AT)**

(72) Inventor: **Eibl, Johann, Dr.
Gustav Tschermak-G. 2
A-1180 Vienna (AT)**
Inventor: **Schwarz, Hans-Peter
Schindlergasse 32
A-1180 Vienna (AT)**
Inventor: **Siekmann, Jürgen, Dr.
Gerasdorfer Strasse 153/209
A-1210 Vienna (AT)**
Inventor: **Turecek, Peter, Dr.
Hutweidengasse 41
A-1190 Vienna (AT)**

(74) Representative: **Kolb, Helga, Dr. Dipl.-Chem. et
al
Hoffmann, Eitle & Partner,
Patent-und Rechtsanwälte,
Arabellastrasse 4
D-81925 München (DE)**

(54) **Stable preparation for the treatment of blood coagulation disorders, comprising an activated coagulation factor and lipid vesicles.**

(57) The invention relates to a stable preparation which comprises a protein that is bound in and/or on lipid vesicles and that was treated for the inactivation of potentially present viruses.

Further, the invention relates to methods for the production of a stable preparation for the treatment of blood coagulation disorders, wherein a protein is bound in and/or on lipid vesicles, and the method comprises a step in which the protein lipid complex is subjected to a treatment for the inactivation of potentially present viruses.

EP 0 680 764 A2

The invention relates to a stable preparation which comprises a protein, wherein the preparation is subjected to a method for virus inactivation.

As a protein, a coagulation active protein is suitable above all, whereby the preparation is suitable for the treatment of blood coagulation disorders. Also, proteins and/or polypeptides with antigenic determinants are possible. An immunologically effective protein has above all an importance as an immunizing agent. In addition, diverse enzymes and/or enzyme inhibitors are understood as proteins as well as other pharmacologically effective proteins.

Coagulation disorders of blood occur when for example the coagulation time deviates from the standard value due to the lack of a coagulation component. The disorders can be inherited as well as acquired through disease conditions.

Hemophilia A is one of the most frequently occurring inherited coagulation disorders. Patients with hemophilia A are prone to frequent hemorrhages as a result of a lack of Factor VIII which can be treated with Factor VIII concentrates. However, in about 15% of the patients the occurrence of Factor VIII neutralizing antibodies, so-called inhibitors, results whereby a therapy with Factor VIII concentrates is hardly possible.

For the treatment of hemophilia A inhibitor patients, complex mixtures of coagulation factors have proven themselves. The activated prothrombin complex FEIBA® (Immuno) has a "factor eight inhibitor bypassing activity".

In lieu of Factor VIII concentrates, it was also attempted to treat hemophilia A with a mixture of coagulation factor Xa and phospholipids. In US 4,610,880, the treatment of hemophilic dogs with a mixture of Factor Xa and phospholipid vesicles is described. This mixture was not stable, for which reason a Factor Xa-containing solution and a suspension of phospholipid vesicles had to be freshly mixed immediately before use. It is stressed that the mixture ratio must be selected in such a way that hemostasis is achieved but thromboses are not caused.

It is known that phospholipid vesicles in combination with Factor Xa increase the danger of thrombosis. On the basis of an in vivo stasis model with rabbits in Blood 59, 401-407 (1982), the increased thrombogenicity of Factor Xa is described when this was tested together with synthetic phospholipids (phosphatidyl choline/phosphatidyl serine lipid vesicles, PCPS vesicles). The danger of thrombosis after administration of prothrombin complex concentrates is also attributed to the combination of coagulation active phospholipids and activated coagulation factors.

It is known from US 4,348,384 to incorporate the coagulation factors Factor VIII and IX in liposomes. Therewith, a preparation which can be orally or intestinally administered for the treatment of hemophilia A or B is made available. Here, the liposomes have a size of 1 µm and protect the Factor VIII or Factor IX from a premature digestion. The administration of these preparations holds no danger of thrombosis because it does not occur intravenously.

The instability of activated coagulation factors in storage is known. A method is known from Thrombosis Research 22, 213-220 (1981) which permits an as stable as possible beta-Factor Xa preparation to be obtained. According to this method, beta-Factor Xa is stored refrigerated in 50% glycerin at pH 7.2 in 0.03 M imidazol buffer. However, a further conversion to inactive fragments also occurs in this method even at 4 °C. In a similar manner, the instability of Factor Xa in stabilizing glycerol-water mixtures is also described in J. Biol. Chem. 248, 7729-7741 (1973).

The object of the invention is to make available a stable preparation for the treatment of blood coagulation disorders comprising a coagulation active protein.

A further object of the invention is to make available a stable preparation which comprises a protein and is subjected to a treatment for inactivation of potentially existing viruses.

The above named objects are solved according to the invention through a stable preparation comprising a complex of suitable proteins and lipids in vesicular form. It has been demonstrated that the stability of an activated coagulation factor is increased in an unexpected manner through the association of an activated coagulation factor such as Factor Xa on a phospholipid in vesicular form, i.e. through the binding in and on a lipid vesicle. Not only an exceptionally good storage stability in solution or frozen and/or lyophilized form is achieved therewith. It can even be demonstrated that an activated coagulation factor is protected from physical inactivation such as for example in a lyophilization process or in a treatment for the purpose of virus inactivation. The in vivo activity of the activated coagulation factor is also maintained, which is also a measure for the stability of the preparation.

Due to the unexpected high stability of the preparations according to the invention, it is possible to subject a protein, for example a blood factor of intrinsic and/or extrinsic coagulation respectively and/or a cofactor and/or an inhibitor of blood coagulation and/or a coagulation active lipoprotein, such as for example Lp(a), or an antigen, especially a viral antigen, in the form bound in or on lipid vesicles to a treatment for

virus inactivation, preferably a chemical and/or physical treatment and especially a heat treatment. Thereby, the biological activity and/or antigenicity remains largely maintained.

A preparation according to the invention is considered as stable when the activity of the protein is retained to more than 40%, preferably more than 50% through measures such as the lyophilization and reconstitution in the absence of customary stabilizers in unbuffered solution and/or when more than 40% of the activity of the active protein, preferably more than 50%, most preferably more than 60%, is retained by the storage of an aqueous solution of the preparation in the absence of customary stabilizers at 22 °C after 20 hours.

The term "coagulation active protein" comprises, according to the invention, an activated or non-activated blood factor of intrinsic and/or extrinsic coagulation, a cofactor of blood coagulation and a coagulation active lipoprotein, such as for example Lp(a), as well as combinations thereof.

The advantage of an antigen in a preparation according to the invention is above all that, simultaneously with an increased stability of the preparation, the lipid components have an adjuvant effect. Thereby, the preparation according to the invention is especially suitable as an immunizing agent. The antigen is for example a recombinant protein and/or polypeptide respectively recombinantly produced. Thereby, depending on the type of the cell culture, a contamination with infectious agents can occur. Through the preparation according to the invention, which is considered as safe in view of a transmission of infectious agents, an infection is however effectively avoided.

As determined by the exceptionally high stability, which was also established in vivo after infusion of the preparation according to the invention, the preparation is exceptionally suitable for the treatment of patients, especially for the treatment of blood coagulation disorders.

Based on its structure, which is characterized by a protein and a lipid component, a complex of a thrombocytic coagulation active substance, i.e. for example a blood coagulation factor that is naturally present in thrombocytes, is suitable according to the invention as a thrombocyte substitute alone or in combination with further coagulation active substances and a lipid vesicle, and, therewith, for the treatment of blood coagulation disorders which are connected with a lack of activated thrombocytes. For one thing, the complex according to the invention imitates the form of a blood platelet, for another functional similarities can be established, such as for example aggregation capability in a protein solution or in the presence of calcium ions or the adsorption on structural proteins of the collagen type. Therefore, the preparation according to the invention can fulfill functions of thrombocytes.

An in vitro test for the determination of the coagulation time of hemophilia A inhibitor plasma serves for the assessment of the effectiveness of the preparation according to the invention for the treatment of hemophilia A inhibitor patients. An effective preparation shortens the coagulation time by at least 50% and preferably to the coagulation time of normal plasma.

It has been surprisingly demonstrated that, through the association of coagulation factor Xa and phospholipid vesicles, the production of a preparation with good storage stability and a high half-life in a Factor VIII inhibitor plasma is possible. The preparation according to the invention is preferably stored in lyophilized form. The lyophilized preparation can be reconstituted to a solution which comprises the complex in vesicular structure. An addition of carbohydrates such as mono- or disaccharide, approximately in an amount of 3 - 20% (w/w) is advantageous. The stability of the complex according to the invention is further increased in the presence of small amounts of calcium ions. It has been further demonstrated that the complex of coagulation factors and lipid vesicles is not thrombogenic in the pharmaceutical preparations for the treatment of coagulation disorders.

As an active component, the preparation according to the invention comprises a vitamin K dependent protein, for example a factor selected from the group of Factors IIa, VIIa, IXa and Xa, preferably Xa β . Furthermore, additional proteins can be included, such as for example the Factors II, VII, IX, X, protein C, activated protein C, protein S and protein Z. It has emerged that the combination of Factor Xa with at least one of the named proteins is particularly effective. An additional amount of Factor VIII, activated Factor VIII, and/or vWF, Factor V and/or activated Factor V is advantageous. A further embodiment of the preparation according to the invention comprises a complex of an activated blood coagulation factor and lipid vesicles and additionally "tissue factor" or fragments of "tissue factor".

The proteins included in the complex are preferably human and isolated from a plasma fraction or of recombinant origin. Based on a potential infectiousness, a treatment for the inactivation of infectious agents, such as for example viruses or prions, is appropriate. It is recommended to undertake a heat treatment of the substances or the complex of the proteins and lipid vesicles. The treatment can ensue for example according to EP 0 159 311.

According to a preferred embodiment, a complex of a highly purified Factor Xa (at least 100 U/mg protein), which is free of infectious agents by a treatment for virus inactivation, with phospholipid vesicles is

made available.

Based on its stability, the preparation is to be made available as a liquid preparation without anything further, i.e. without addition of the commonly used stabilizers such as carbohydrates (saccharose) or protease inhibitors (aprotinin).

However, the preparation can in addition also be stored in liquid deep-frozen form or in lyophilized form wherein a polysaccharide amount is advantageous in water removal.

According to the invention, phospholipids in vesicular form can be used as lipid vesicles. The phospholipids can be synthetic or of natural origin. Thereby, a standardized mixture of chemically pure phospholipids is advantageous.

The lipid vesicles can be prepared in various ways such as by sonification of phospholipid dispersions (Barenholz et al., *Biochemistry* 16, 2806-2810, 1977), the reversed phase evaporation technique (F. Szoka and D. Paphadjopoulos, *Proc. Natl. Acad. Sci. USA* 75, 4194-4198, 1978), the ethanol injection method (S. Batzri and E.D. Korn, *Biochem. Biophys. Acta* 298, 1015-1019, 1973) or also the removal of detergents from mixed micelles by dialysis methods (O. Zumbuehl and H.G. Weder, *Biochem. Biophys. Acta* 640, 252-262, 1981). Furthermore, the extrusion method according to Olson et al. (*Biochem. Biophys. Acta* 557, 9-23, 1979) can also be used. A dispersion of multilamellar vesicles is prepared by hydration of a lipid film which can be obtained preferably by evaporation of solutions of the lipids in organic solvents with the aid of a rotary evaporator. This dispersion is subsequently pressed through two stacked polycarbonate filters with nitrogen pressure.

This method should be carried out at a temperature at which the phospholipids are found in a liquid-crystalline state (at least 5°C, preferably at least 10°C above the phase transition temperature, T_M). Optionally, a repeated freeze-thaw cycle is introduced before the extrusion (M.J. Hope et al., *Biochem. Biophys. Acta* 812, 55-65, 1985). Furthermore, an embodiment has proven to be particularly advantageous. Here, the dispersion of multilamellar vesicles is produced as outlined above, lyophilized, and reconstituted again with water. Thereby, the subsequent extrusion can be carried out particularly easily and also with very high lipid concentrations (for example, 100 mg lipid/ml).

For the production of the complex according to the invention, the described lyophilization step can be carried out after mixing of the vesicle dispersion with the protein, here preferably coagulation factor Xa. After reconstitution, the complex according to the invention can be further extruded. A further possibility for the production of the complex according to the invention is the mixing of the components in the presence of a tenside and the subsequent removal of the tenside, for example by dialysis. A direct hydration of the lipid film with a solution of proteins is also particularly advantageous, wherein optionally lyophilization and reconstitution can be done, whereby the extrusion is made easier. Optionally, in the last variation, the additional lyophilization step can be dispensed with. A direct mixing of the prepared vesicles with the protein preferably in the presence of charged particles or under conditions which permit the incorporation of the protein in and on the lipid vesicles (concentration, selection of proteins with hydrophobic properties, etc.) is also conceivable.

For the extrusion process, particularly filters with a diameter of about 30 to 1000 nm can be used. The typical size of the lipid vesicles (determined by the method of dynamic light scattering) is in the order of 30 to 900 nm, preferably 100 to 500 nm.

The stability of the produced complex is improved when charged particles, for example calcium ions, preferably 1-10 mM, are present. Based on the binding of the coagulation proteins in and on the lipid vesicles, a purification and isolation of the complex per se is possible. This can occur by gel filtration or ultracentrifugation.

The glycerophospholipids phosphatidyl serine, phosphatidic acid, phosphatidyl glycerol, diphosphatidyl glycerol (cardiolipin) and also phosphatidyl ethanolamine which can exhibit a negative charge are among the suitable phospholipids and are used preferably alone or also combined in a mixture with neutral phospholipids such as phosphatidyl choline and sphingomyelin, wherein the binding by addition of calcium ions is favored. As anionic components, the use of other lipids, such as for example sulfatides or also dicetylphosphate, phosphatidyl methanol, phosphatidyl- β -lactate and oleic acid is also possible. However, the exclusive use of neutral phospholipids such as phosphatidyl choline or sphingomyelin is also conceivable. However, substances obtained through chemical modification such as the lipid derivatives of the polyethylene glycols or substances from the classes of lysophospholipids or glycolipids are also useable. The use of lipids which can exhibit a positive charge, such as for example stearylamine, dimethyldioctadecyl ammonium bromide (DDAB), or also cationic lipids of the type of the 1,2-diacyl-3-trimethyl ammonium propanes (TAP) or the 1,2-diacyl-3-dimethyl ammonium propanes (DAP) is also conceivable.

A preferred embodiment comprises the use of lipids with anti-viral effect, for example alkyl-phospholipids (see EP 0 506 704 B1).

For stabilization, the phospholipid vesicles can comprise up to 80 Mol% cholesterol as an additional component. It is to be emphasized that all specified lipids can also be used as a sole component, optionally also under addition of cholesterol.

For the formation of the complex the lipid vesicles can be found in the gel state or in the liquid-crystalline state, however, the use of systems in the liquid-crystalline state is more favorable. In a version of the invention, lipid vesicles comprised of 1,2-dioleoyl-sn-glycero-3-phosphocholine and 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoserine are used (for example in a ratio of PC/PS = 80/20 (w/w), see Example 1) which are liquid-crystalline at room temperature. The use of phospholipids with unsaturated side chains can, however, lead to a reduced stability by virtue of oxidation. Therefore, for the chemical stability of the vesicles, the use of lipids with fatty acid groups such as those of myristic acid, palmitic acid (Examples 2 and 3) or stearic acid, which are however normally found in the gel state, is preferable. The use of these lipids has been proven as particularly advantageous, however, in mixtures with cholesterol.

The invention is more closely illustrated through the following Examples.

15 **Production of phospholipid vesicles**

Example 1: Production of multilamellar vesicles

In a 100 ml flask, 40.0 mg 1,2-dioleoyl-sn-glycero-3-phosphocholine and 10.0 mg 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoserine (Avanti Polar Lipids) are dissolved in 5 ml chloroform and evaporated with the aid of a rotary evaporator under reduced pressure and a temperature of 30 °C. After complete removal of the solvent, a vacuum was still held over 30 min at 30 mbar and subsequently dried over a time period of six hours in high-vacuum at 0.1 mbar. The phospholipid film was subsequently hydrated by addition of 5 ml of buffer (20 mM Tris, 150 mM NaCl, pH 7.4) and gentle shaking over one hour at room temperature.

Example 2

After formation of a dispersion of multilamellar vesicles according to Example 1, this was extruded 10 times with N₂ pressure through two stacked 100 nm polycarbonate filters laid on top of each other (10 ml thermobarrel extruder, Lipex Biomembranes Inc., Vancouver, Canada). The determination of the particle size of the vesicles produced in this way with the aid of dynamic light scattering (Malvern Zetasizer 4) resulted in a average diameter of about 100 nm.

Example 3

From 35.0 mg 1,2-dimyristoyl-sn-glycero-3-phosphocholine (Nattermann Phospholipid GmbH) and 15.0 mg 1,2-dimyristoyl-sn-glycero-3-phosphoserine (Avanti Polar Lipids) in 5 ml chloroform/methanol mixture (2:1, v/v), a phospholipid film was produced as described in Example 1. After addition of 5 ml of buffer (20 mM Tris, 150 mM NaCl, pH 7.4), the film was hydrated at 50 °C by using a rotary evaporator and extruded at 50 °C as described under Example 2.

Example 4

From a mixture of 35.0 mg 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (Nattermann Phospholipid GmbH) and 15.0 mg 1,2-dimyristoyl-sn-glycero-3-phosphoserine (Avanti Polar Lipids), phospholipid vesicles were produced as described in Example 3. The hydration of the film and the extrusion also occurred at 50 °C.

Example 5

From a mixture of 40.0 mg 1,2-dioleoyl-sn-glycero-3-phosphocholine (Sigma) and 10.0 mg 1,2-dioleoyl-sn-glycero-3-phosphoglycerol or 17.5 mg cardiolipin from bovine heart (Sigma), phospholipid vesicles were produced as described under Example 1 and 2.

Example 6

From a mixture of 40.0 mg 1,2-dioleoyl-sn-glycero-3-phosphocholine and 10.0 mg 1,2-dioleoyl-sn-glycero-3-phosphate (Sigma), phospholipid vesicles were produced as described under Example 1 and 2.

Example 7

From a mixture of 35.0 mg 1,2-dioleoyl-sn-glycero-3-phosphocholine, 10.0 mg 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoserine (Avanti Polar Lipids) and 5 mg phosphatidyl inositol from Soya beans (Sigma),
 5 phospholipid vesicles were produced as described under Example 1 and 2.

Example 8

5 to 80 Mol% cholesterol (Sigma) was added to the mixtures of phosphatidyl choline, phosphatidyl
 10 serine, phosphatidyl glycerol, phosphatidic acid and/or phosphatidyl inositol described in Examples 1 to 7
 and extruded as described under the respective Examples.

Example 9

15 After hydration of the phospholipid film, the dispersions from Examples 1-8 were shock frozen (liquid
 N_2) and thawed at 37 °C (Examples 2, 5-8) and/or 50 °C (Examples 3 and 4). This process was repeated
 four times. The multilamellar vesicles produced in this way were extruded as described in the respective
 Examples. The vesicle preparations resulting therefrom had an average diameter of 100 nm.

Example 10

As described in the Examples 1-9, multilamellar vesicles were produced through hydration of
 phospholipid films. The dispersions obtained in this way were frozen at -80 °C and lyophilized. After
 reconstitution of the lyophilizates with the corresponding volumes of H_2O , extrusion was done as described
 25 under the respective Examples.

Example 11

The production of phospholipid vesicles according to Examples 1-10 was carried out in 150 mM NaCl
 30 instead of Tris buffer. The size determination of the vesicles with the aid of dynamic light scattering resulted
 in a nearly identical size distribution in comparison to vesicles that were produced with buffered solution.

Example 12: 1,2-dioleoyl-sn-glycero-3-phosphocholine

35 A mixture of 8.0 mg 1,2-dioleoyl-sn-glycero-3-phosphocholine (Avanti Polar Lipids), 2.0 mg phosphatidyl
 inositol from Soya beans (Sigma) and 10.0 mg sodium cholate (Sigma) was added in a 50 ml round flask
 and dissolved in 10 ml of a chloroform/methanol mixture (1:1, v/v). Then, this was evaporated in a rotary
 evaporator at a bath temperature of 20 °C under reduced pressure until dry, taken up in 10 ml absolute
 40 methanol and evaporated again. The film produced in this way was taken up under shaking with 2 ml buffer
 (20 mM Tris, 150 mM NaCl, pH 7.4). For the production of the vesicles, the clear solution was subsequently
 dialyzed 24 h against the same buffer at 4 °C. The determination of the particle size with the aid of dynamic
 light scattering resulted in an average diameter of about 70 nm.

Lyophilization of phospholipid vesicles in the presence of saccharose

45

Example 13

Saccharose to a concentration of 3 - 20% (w/v) was added to the vesicles produced corresponding to
 the Examples 1-12. Subsequently, this was frozen at -80 °C and lyophilized. After reconstitution of the
 50 lyophilizate, the determination of the size with the aid of dynamic light scattering resulted in an average
 diameter of about 100 nm. The electron microscopic examination of this solution resulted in a size
 distribution of about 80 to 130 nm according to visualization with the negative staining method.

55

Production of a complex of Factor Xa and phospholipid vesicles**Example 14: colyophilization**

5 Factor Xa was produced as follows:

A solution of the prothrombin complex preparation (corresponding to 50,000 U Factor X/l) produced according to the methods of Brummelhuis (Methods of Plasma Protein Fractionation, J.M. Curling (editor), pp 117, Acad. Press., 1980) and heat treated according to the method of EP 0 159 311, was treated in the presence of 12 % (v/v) Tween®80 one hour at pH 7.0 and room temperature. Subsequently, this was
 10 diluted, mixed with trisodium citrate dihydrate (7.0 g/l), and the proteins of the prothrombin complex precipitated at pH 7.0 by addition of 400 ml 1 M barium chloride solution. The precipitate was washed and resuspended in a 25 % (w/v) ammonium sulfate solution containing 50 mM benzamidine HCl. The non-dissolved matter was separated and the solution was brought to 80% saturation with ammonium sulfate in order to precipitate the proteins again. The precipitate was dissolved in buffer and rebuffed
 15 chromatographically over Sephadex®-G25 against 25 mM trisodium citrate buffer containing 100 mM NaCl and 1 mM benzamidine HCl, pH 6.0.

The protein containing fractions were further purified by means of a DEAE Sepharose® fast flow column. By increasing the sodium chloride concentration in the citrate buffer (25 mM, pH 6.0), Factor X was eluted separated from the other prothrombin complex proteins. The obtained fraction was rebuffed against
 20 20 mM Tris HCl buffer containing 150 mM sodium chloride and 2 mM calcium chloride (pH 7.4). Subsequently, this was mixed with 0.14 mg RVV (Russel's viper venom, a protease from Vipera russellii, Pentapharm) per 100 U FX and stirred one hour at room temperature. The generated Factor Xa was then purified chromatographically with a benzamidine Sepharose® column and concentrated through ammonium sulfate precipitation and subsequently chromatographed over Sephadex®-G25 for the removal of the salt.
 25 The produced Factor Xa preparation had a specific activity of at least 100 U Factor Xa/mg protein.

For the production of the complex, a vesicle preparation produced according to the Examples 1-12 was mixed, with or without 2.5 mM calcium chloride, to the Factor Xa fraction. The solution was lyophilized.

After reconstitution, the effectiveness of the preparation in the FEIBA test (test description see AT 350 726) was tested. As a comparison for this, the FEIB activity of the non-complexed Factor Xa was
 30 determined. Before the lyophilization, the FEIB activity amounted to 1700 U/ml in both cases. After the lyophilization and reconstitution, 1650 U/ml was recovered in the case of the Factor Xa/PCPS complex. The loss of FEIB activity was much larger in the case of non-complexed Factor Xa. Only 1220 U/ml was recovered.

35 Example 15: Extrusion of a colyophilizate

A colyophilizate produced according to Example 14 comprising 1000 U Factor Xa as well as 50.0 mg of a phospholipid preparation produced according to Example 6 of 40.0 mg 1,2-dioleoyl-sn-glycero-3-phosphocholine and 10.0 mg 1,2-dioleoyl-sn-glycero-3-phosphate was taken up with water in the original
 40 volume and extruded twice at a temperature of 20 °C through two stacked 400 nm polycarbonate filters (Lipex Biomembranes, Inc.). The determination of the particle size with the aid of dynamic light scattering (Malvern Zetasizer 4) resulted in an average diameter of about 450 nm.

Example 16: coextrusion

45 As described under Example 1, a film was produced from a phospholipid mixture. This was hydrated with a Factor Xa containing solution from Example 14 with or without 2.5 mM calcium chloride. After hydration of the film, it was proceeded as in the Examples 2, 9 and 10 and extruded. Thereby, Factor Xa was complexed with phospholipid vesicles. Saccharose in a concentration of 3 - 20 % (w/v) could be added
 50 to the complex produced in this way in order to subsequently lyophilize.

Production of a complex of activated protein C, Factor Xa and phospholipid vesicles**Example 17**

55 A film was prepared as described under Example 1 from a phospholipid mixture. This was hydrated with the Factor Xa containing fraction from Example 14 as well as with activated protein C in a 20 mM Tris HCl buffer containing 150 mM NaCl and 5 mM CaCl₂, pH 7.4. The mixture containing 10 U FXa/ml, 10 U

APC/ml and 10 mg phospholipid/ml was lyophilized.

Purification of the complex of Factor Xa and PCPS vesicles

5 Example 18

A preparation containing Factor Xa and PCPS vesicles was produced according to the method described in the Examples 14 or 16, wherein saccharose in a final concentration of 5 % (w/v) was added to the Factor Xa phospholipid preparation. The lyophilizate was dissolved in water in such a way that the prepared solution contained 2.4 U Factor Xa/ml and 1 mg/ml PCPS vesicles. A column packed with Superose 6 HR 10/30, Pharmacia, was equilibrated with a buffer (20 mM Tris HCl containing 150 mM NaCl, 0.1 % albumin, 0.01 % Tween 20, 1 mM CaCl₂, pH 7.4). 500 µl of the PCPS vesicles and Factor Xa containing preparation were chromatographed over the column with a flow rate of 0.4 ml/minute.

In the eluate stream, the UV absorption at 254 nm was measured. The fraction of the exclusion volume containing a complex of PCPS vesicles and Factor Xa was collected.

Example 19

A lyophilizate produced according to Example 14 was reconstituted with water such that the finished solution contained 5 U Factor Xa and 5 mg PCPS per ml. 0.5 ml of this solution were mixed with 1 ml of a 30 % (w/v) ficoll 400 solution (Pharmacia, in 150 mM sodium chloride solution). This mixture was placed in an ultracentrifugation tube and overlaid with 3 ml of 10 % ficoll solution. Finally, this was overlaid with 150 ml NaCl solution and subsequently centrifuged by use of a swing out rotor for 30 minutes at 100,000 g and room temperature.

The layer containing the purified Factor Xa/PCPS complex with lower density than the aqueous solution could be separated as a supernatant.

Example 20

A complex comprising activated protein C, Factor Xa and phospholipid vesicles was produced according to Example 17. The lyophilized complex was reconstituted with water and concentrated to a third of the starting volume by centrifugation over ultrafree-MC filter units, exclusion limit 100,000 Daltons (polysulphone membranes, Millipore), 30 minutes at 4000 rpm. The retained material obtained in this way contained the purified complex of activated protein C, Factor Xa and phospholipid vesicles.

Stability of the complex of protein and phospholipids

Example 21: Stability of a Factor Xa/PCPS vesicle complex in the lyophilization process

PCPS vesicles were produced as described in Examples 1 and 2. These were subsequently mixed with Factor Xa, which was produced as described in Example 14, such that the preparation comprised 0.5 mg phospholipid vesicles per ml and 47 U Factor Xa in an aqueous solution of 150 mM/l NaCl and 5 mM/l CaCl₂. The complex was lyophilized. As a comparison for this, Factor Xa was lyophilized in the same concentration, however without PCPS vesicles. Subsequently, according to German patent application P4325872.7, the Factor Xa activity was determined in the lyophilized preparations after reconstitution with distilled water in the starting volumes and compared with the Factor Xa amount before the lyophilization (see Table).

	Factor Xa activity (%)	
	before lyo	after lyo
PCPS/ FXa-complex	100	51
Factor Xa	100	20

The Table shows the stabilizing influence of vesicular phospholipid on Factor Xa in the copreparation.

Example 22: Stability of a Factor Xa/PCPS vesicle complex in solution

A complex of PCPS vesicles and Factor Xa was produced as described in Example 21 and lyophilized. After the reconstitution, the solution was stored at 22 °C over 20 hours. Subsequently, the activity of Factor Xa was determined. The following Table shows the stability of Factor Xa in the PCPS vesicle complex in comparison to non-complexed Factor Xa.

	Factor Xa activity (%)	
	0 h	20 h
PCPS/ FXa-complex	100	67
Factor Xa	100	34

Example 23: Stability of a protein C/Factor Xa/PCPS vesicle complex in the lyophilization

A complex comprising protein C, Factor Xa and PCPS vesicles was produced analogously to Example 17 and lyophilized. The preparation contained 0.5 mg phospholipid vesicles/ml, 10 U protein C/ml and 47 U Factor Xa/ml in an aqueous solution of 150 mM NaCl. As a comparison for this, protein C and Factor Xa was lyophilized at the same concentrations, however without PCPS vesicles. Subsequently, the Factor Xa activity was determined as in Example 21 and the protein C activity was determined by use of a chromogenic test (Immunochrom PC, Immuno) in the lyophilized preparations after reconstitution with distilled water in the starting volumes and compared with the respective amount of Factor Xa and protein C before the lyophilization (see Table).

	protein C activity (%)		Factor Xa activity (%)	
	before lyo	after lyo	before lyo	after lyo
PCPS/protein C/FXa complex	100	89	100	84
protein C	100	41	-	-
Factor Xa	-	-	100	23

The Table shows the stabilizing influence of phospholipid vesicles on protein C and Factor Xa in the complex.

Example 24: Stability of a protein C/FXa/PCPS vesicle complex in solution

A complex of protein C, Factor Xa and PCPS vesicles and Factor Xa was produced as described in Example 23 and lyophilized. After the reconstitution, the solution was stored at 22 °C over 20 hours. Subsequently, the activity of protein C and Factor Xa was determined. The following Table shows the increased stability of protein C and Factor Xa in the PCPS vesicle complex in comparison to the non-complexed factors.

	protein C activity (%)		Factor Xa activity (%)	
	0 h	20 h	0 h	20 h
PCPS/protein C/FXa-complex	100	100	100	88
protein C	100	65	-	-
Factor Xa	-	-	100	9

In vitro characterization of the protein/phospholipid complexes.**Example 25**

For the testing of the effectiveness of the preparation according to the invention, the following test system was used.

100 μ l of FVIII inhibitor plasma (45 Bethesda Units/ml) were mixed in a coagulometer tube with 100 μ l 20 mM Tris HCl buffer containing 150 mM NaCl, pH 7.4 (TBS) and recalcified with a further 100 μ l of a 0.025 M calcium chloride solution. Immediately after addition of the sample to be analyzed (100 μ l), the coagulation time of the mixture was determined by use of a coagulometer (Schnitger/Gross) at 37 °C.

When pure TBS buffer was employed as a sample in the represented test substance, the coagulation time amounted to 820 seconds. In comparison to this, the coagulation time of normal plasma in this test system was about 350 seconds.

A preparation produced according to Example 16 was tested at a concentration of 0.88 mU Factor Xa and 0.3 μ g phospholipid/ml. As an additive, either 0.1 U FEIBA produced according to the method of AT 368 883 or 0.1 U recombinant Factor VIIa (Novo Nordisk) were tested. Also, a mixture of the FEIBA preparation (0.01; 0.1; 1.0 U/ml) or of Factor X (0.01; 0.1; 1.0 U/ml) produced according to the method from Example 14 with 0.3 μ g/ml phospholipid vesicles produced according to the method from Example 1 and 2 were tested. The influence of the tested substances on the coagulation time is given in the following Table.

Shortening of the coagulation time by a preparation containing the complex of coagulation protein and phospholipid vesicles.

test substance	concentration (mU/ml)				coagulation time (s)
	FXa	FEIBA	FVIIa	FX	
Factor Xa/PCPS	0.88	-	-	-	125
Factor Xa/FEIBA/PCPS	0.88	100	-	-	82
Factor Xa/Factor VIIa/PCPS	0.88	-	100	-	131
FEIBA/PCPS	-	10	-	-	409
		100			167
		1000			120
Factor X/PCPS	-	-	-	10	814
				100	425
				1000	336

Biological in vitro activity of various phospholipid vesicle types**Example 26**

Phospholipid vesicles comprising mixtures of various phospholipid types in different composition were produced as described in the Examples and processed to copreparations with Factor Xa. The FXa phospholipid vesicle complex resulting therefrom was examined as described in Example 25 on the coagulation shortening effect of FVIII inhibitor plasma. The results (coagulation times) are to be taken from the following Table:

EP 0 680 764 A2

vesicle type	composition w/w %	conc./ml in test substance		coagulation time (s)
		mU FXa	µg lipid	
DOPC	80	1.375	6.25	62
POPS	20			
DOPC	70	1.375	6.25	65
POPS	20			
CHOL	10			
DOPC	60	1.375	6.25	57
POPS	20			
CHOL	20			
DOPC	50	1.375	6.25	56
POPS	20			
CHOL	30			
DOPC	40	1.375	6.25	69
POPS	20			
CHOL	40			
DOPC	30	1.375	6.25	58
POPS	20			
CHOL	50			
POPS	95	1.375	6.25	73
PI	5			
POPS	90	1.375	6.25	77
PI	10			
POPS	80	1.375	6.25	71
PI	20			
DOPC	80	1.375	6.25	135
PI	20			
DOPC	80	1.375	6.25	66
POPS	15			
PI	5			
DOPC	80	1.375	6.25	89
POPS	10			
PI	10			
DOPC	80	1.375	6.25	110
POPS	5			
PI	15			

	vesicle type	composition w/w %	conc./ml in test substance		coagulation time (s)
			mU FXa	µg lipid	
5	DOPC	75	1.375	6.25	75
	POPS	20			
	PI	5			
	DOPC	70	1.375	6.25	80
	POPS	20			
	PI	10			
10	DOPC	60	1.375	6.25	72
	POPS	20			
	PI	20			
	DOPC	60	1.375	6.25	85
	POPS	10			
15	PI	10			
	CHOL	20			
	DOPC	50	1.375	6.25	82
	POPS	20			
	PI	10			
20	CHOL	10			
	CLP	100	1.16	5.63	76
	DOPC	70	1.16	6.88	69
	CLP	30			
	DMPC	70	1.16	8.45	91
25	DMPS	30			
	DPPC	70	1.16	8.45	123
	DMPS	30			
	DMPC	50	1.16	8.45	86
	DMPS	30			
30	CHOL	20			
	DMPC	30	1.16	8.45	67
	DMPS	30			
	CHOL	40			
	DPPC	50	1.16	8.45	100
35	DMPS	30			
	CHOL	20			
	DPPC	30	1.16	8.45	154
	DMPS	30			
	CHOL	40			
40	DOPG	80	1.25	10	77
	DOPC	20			
	DOPA	80	1.25	10	67
	DOPC	20			
45	buffer background		0	0	820

Legend:

5	DOPC	=	1,2-dioleoyl-sn-glycero-3-phosphocholine
	POPS	=	1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoserine
	CHOL	=	cholesterol
10	CLP	=	diphosphatidyl glycerin (cardiolipin)
	PI	=	phosphatidyl inositol
	DMPC	=	1,2-dimyristoyl-sn-glycero-3-phosphocholine
15	DMPS	=	1,2-dimyristoyl-sn-glycero-3-phosphoserine
	DPPC	=	1,2-dipalmitoyl-sn-glycero-3-phosphocholine
	DOPG	=	1,2-dioleoyl-sn-glycero-3-phosphoglycerin
20	DOPA	=	1,2-dioleoyl-sn-glycero-3-phosphoric acid

Example 27

25 The effectiveness of a complex produced according to Example 14 with 20 mU Factor Xa and 4.96 μ g PCPS/ml, and various dilutions thereof, in Factor VIII inhibitor plasma was determined in the following coagulation test immediately after mixing and after an hour incubation at 37°C:

200 μ l of sample were mixed in a coagulometer tube with 100 μ l 20 mM Tris HCl buffer containing 150 mM NaCl, pH 7.4 (TBS) and recalcified with a further 100 μ l of a 0.025 M calcium chloride solution.
 30 Immediately after addition of this solution, the coagulation time of the mixture was measured by use of a coagulometer (Schnitger/Gross) at 37°C.

When a 1 + 1 mixture of FVIII inhibitor plasma (45 Bethesda Units/ml) and TBS buffer was employed as a sample in the represented test substance, the coagulation time amounted to more than 500 seconds.

As a comparison, pure Factor Xa produced according to the method from Example 14 was tested. The
 35 results are summarized in the following Table.

Coagulation time of Factor Xa/PCPS and Factor Xa before and after incubation with Factor VIII inhibitor plasma

40

45

50

55

5	dilution of the stock solution	coagulation time (s)			
		Factor Xa		Factor Xa/PCPS	
		0 h	1 h	0 h	1 h
10	undiluted	77	161	35	69
	1:2	95	236	48	111
	1:4	117	349	65	185
15	1:8	152	>500	88	314
	1:16	224	>500	145	400
	1:32	314	>500	200	>500
	1:64	441	>500	296	>500
20	1:128	>500	>500	402	>500

25 The complex of Factor Xa/PCPS according to the invention has a higher stability than non-complexed Factor Xa. In low concentrations, Factor Xa/PCPS also leads to a shortening of the coagulation time after one hour incubation in Factor VIII inhibitor plasma, whereas non-complexed Factor Xa had already lost its coagulation time shortening activity.

30 In vivo characterization of the protein/phospholipid complexes

30 Example 28: Test on thrombogenicity

35 A FEIBA preparation produced according to the method of AT 368 883 was tested for thrombogenicity as a complex with PCPS vesicles and in non-complexed form in the Wessler-Stasis model. The PCPS vesicles were produced according to the method from Example 1 and 2, wherein vesicles of average sizes of 100 nm and/or 50 nm were obtained through a suitable filter choice. 4 U FEIBA and 60 µg PCPS were used per kg rabbit. The thrombogenicity of the FEIBA preparation was not increased by the complex formation with PCPS vesicles. In all cases, no thrombi formation was registered in the stasis model (score = 0-1).

40 Heat treatment of the protein/phospholipid vesicle complexes

Example 29: Heat treatment of a complex of Factor Xa and PCPS vesicles

45 A lyophilizate of Factor Xa and PCPS produced according to Example 14 or 16 was treated according to the method of EP 159 311 for 10 hours at 60 °C and 1 hour at 80 °C. The Factor Xa activity of the reconstituted solution amounted to more than 90% of the activity before the heat treatment. With the aid of the method of dynamic light scattering (Malvern Zetasizer 4) the preservation of the vesicular structure could be detected.

50 Example 30: Heat treatment of a complex of FEIBA and PCPS vesicles

55 A complex of phospholipid vesicles and FEIBA was produced analogously to Example 14 and lyophilized. After reconstitution, the lyophilizate contained 30 U FEIBA/ml and 3.4 mg PCPS/ml in a buffer of 4 g Na₃citrate•2H₂O/l and 8 g NaCl/l, pH 7.0. A lyophilizate corresponding to this composition was treated according to the method of EP 159 311 for 10 hours at 60 °C and 1 hour at 80 °C. The FEIBA activity in the reconstituted solution after heat treatment amounted to more than 90% of the activity before the heat treatment. With the aid of the method of dynamic light scattering (Malvern Zetasizer 4) the preservation of

the vesicular structure could be detected.

Claims

- 5 1. Stable, virus safe preparation comprising a protein that is bound in and/or on lipid vesicles, characterized in that the preparation was treated for the inactivation of potentially present viruses.
2. Preparation according to claim 1, characterized in that the protein is coagulation active.
- 10 3. Preparation according to claim 1, characterized in that the preparation is treated by chemical and/or physical measures.
4. Preparation according to claim 3, characterized in that the preparation is heat treated.
- 15 5. Preparation according to any one of the claims 1 to 4, characterized in that a blood factor of the intrinsic or extrinsic coagulation is included.
6. Preparation according to claim 5, characterized in that a blood coagulation factor or cofactor in activated form is included.
- 20 7. Preparation according to claim 6, characterized in that the activated coagulation factor is a vitamin K dependent protein.
8. Preparation according to claim 7, characterized in that the activated coagulation factor is selected from the group consisting of Factors IIa, VIIa, IXa and Xa.
- 25 9. Preparation according to claim 8, characterized in that Factor Xa β is included.
10. Preparation according to any one of claims 6 to 9, characterized in that a protein selected from the group consisting of Factor II, VII, IX, X, protein C, activated protein C, protein S and protein Z is additionally included.
- 30 11. Preparation according to any one of claims 6 to 10, characterized in that "tissue factor" or a fragment of "tissue factor" is additionally included.
- 35 12. Preparation according to any one of claims 6 to 11, characterized in that Factor VIII, activated Factor VIII and/or von Willebrand Factor is additionally included.
13. Preparation according to any one of claims 5 to 11, characterized in that Factor V and/or activated Factor V is additionally included.
- 40 14. Preparation according to any one of claims 1 to 4, characterized in that the protein is an enzyme and/or an inhibitor.
- 45 15. Preparation according to any one of claims 1 to 4, characterized in that the protein is a coagulation active lipoprotein.
16. Preparation according to claim 15, characterized in that the lipoprotein is Lp(a).
- 50 17. Preparation according to any one of claims 1 to 4, characterized in that the protein is an antigen.
18. Preparation according to one of the above claims, characterized in that it is suitable for intravenous administration.
- 55 19. Preparation according to one of the above claims, characterized in that the lipid vesicles have a size determined according to the method of dynamic light scattering in the range of 30 to 900 nm, preferably 100 to 500 nm.

20. Preparation according to one of the above claims, characterized in that it does not comprise common stabilizers such as carbohydrates or protease inhibitors.
- 5 21. Preparation according to one of the above claims, characterized in that it is present in liquid-frozen or lyophilized form.
22. Preparation according to one of the above claims, characterized in that the protein with a biological activity of at least 90% with respect to the activity before the inactivation treatment is included.
- 10 23. Method for the production of a preparation according to claim 1, characterized in that a lyophilization step is carried out after mixing a vesicle dispersion with the protein and that the method comprises a step in which the protein bound in and/or on lipid vesicles is subjected to a treatment for virus inactivation.
- 15 24. Method for the production of a preparation according to claim 1, characterized in that a direct hydration of a lipid film occurs with a solution of protein and that the method comprises a step in which the protein bound in and/or on lipid vesicles is subjected to a treatment for virus inactivation.
- 20 25. Method for the production of a preparation according to claim 1, characterized in that the prepared vesicles are mixed directly with protein in order to permit an incorporation of the protein in and on the lipid vesicles and that the method comprises a step in which the protein bound in and/or on lipid vesicles is subjected to a treatment for virus inactivation.

25

30

35

40

45

50

55

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 680 764 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
28.01.1998 Bulletin 1998/05

(51) Int. Cl.⁶: **A61K 38/48**, C12N 9/96,
A61K 47/24

// (A61K38/48, 38:35)

(43) Date of publication A2:
08.11.1995 Bulletin 1995/45

(21) Application number: **95106821.2**

(22) Date of filing: **05.05.1995**

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB IT LI NL SE

(30) Priority: **06.05.1994 DE 4416166**

(71) Applicant:
IMMUNO Aktiengesellschaft
A-1221 Wien (AT)

(72) Inventors:
• **Eibl, Johann, Dr.**
A-1180 Vienna (AT)

- **Schwarz, Hans-Peter**
A-1180 Vienna (AT)
- **Siekmann, Jürgen, Dr.**
A-1210 Vienna (AT)
- **Turecek, Peter, Dr.**
A-1190 Vienna (AT)

(74) Representative:
Kolb, Helga, Dr. Dipl.-Chem. et al
Hoffmann Eitle,
Patent- und Rechtsanwälte,
Arabellastrasse 4
81925 München (DE)

(54) **Stable preparation for the treatment of blood coagulation disorders, comprising an activated coagulation factor and lipid vesicles**

(57) The invention relates to a stable preparation which comprises a protein that is bound in and/or on lipid vesicles and that was treated for the inactivation of potentially present viruses.

Further, the invention relates to methods for the production of a stable preparation for the treatment of blood coagulation disorders, wherein a protein is bound in and/or on lipid vesicles, and the method comprises a step in which the protein lipid complex is subjected to a treatment for the inactivation of potentially present viruses.

EP 0 680 764 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 95 10 6821

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	GB 2 129 685 A (NAT BIOLOG STANDARDS BOARD) * example 26 * ---	1-6,12, 18, 20-23,25	A61K38/48 C12N9/96 A61K47/24 //(A61K38/48, 38:35)
X	WO 91 02532 A (UNIV KINGSTON) * page 11 - page 12 * ---	1-8,14, 17,18, 20-22	
X	WO 90 03808 A (LIPOSOME CO INC) * the whole document * ---	1,3,4, 14,17-23	
X,P	WO 95 04524 A (OPPERBAS HOLDING BV ;BARENHOLZ YECHZKEL (IL); NUR ISRAEL (IL)) * page 11; claims 1-24; figure 1 * ---	1-8,14, 17-25	
E	EP 0 680 763 A (IMMUNO AG) * the whole document * ---	1-25	
A	WO 93 23074 A (OKLAHOMA MED RES FOUND ;MORRISSEY JAMES H (US); PHILIP C COMP (US)) * page 6 * ---	1-25	TECHNICAL FIELDS SEARCHED (Int.Cl.6) A61K C12N
A	EP 0 214 737 A (UNIV KINGSTON) * the whole document * ---	1-25	
A	EP 0 159 311 A (IMMUNO AG) * the whole document * ---	1-25	
A	GB 2 085 729 A (DAINIPPON PHARMACEUTICAL CO) * the whole document * ---	1-25	
		-/--	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 3 December 1997	Examiner Fernandez y Branas, F
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03 82 (P04C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 95 10 6821

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A,D	US 4 348 384 A (HORIKOSHI ISAMU ET AL) * the whole document * -----	1-25	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 3 December 1997	Examiner Fernandez y Branas, F
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			

EPC FORM 1503 03.82 (Pct/C01)

